

POSTER PRESENTATION

Open Access

β -arrestin 2 regulates A β generation and γ -secretase activity in Alzheimer's disease

Amantha Thathiah^{1,2*}, Katrien Horre^{1,2}, An Snellinx^{1,2}, Elke Vandeweyer^{1,2}, Yunhong Huang^{1,2}, Marta Ciesielska^{1,2}, Gardien De Kloe^{1,3}, Sebastian Munck^{1,2}, Bart De Strooper^{1,2}

From Molecular Neurodegeneration: Basic biology and disease pathways
Cannes, France. 10-12 September 2013

Deficits in several neurotransmitter systems are characteristic features of the brains of AD patients. The majority of neurotransmitters communicate information to cells via G protein-coupled receptors (GPCRs) or 7-transmembrane receptors (7TMRs). It has recently been appreciated that a small family of multifunctional GPCR regulatory known as the β -arrestins which play an almost universal role in facilitating traditional GPCR desensitization, are also capable of initiating distinct signals in their own right, conveying receptor subtype-specific signaling events. These signals are often both spatially and temporally distinct, and result in unique cellular and physiological or pathophysiological consequences. As mediators of GPCR desensitization, trafficking and cell signaling, the β -arrestins provide a putative basis to understand GPCR dysfunction in AD. Here, we report that β -arrestin 2 levels are elevated in two independent cohorts of patients with AD. Genetic deletion of *Arrb2* (β -arrestin 2) reduces accumulation of the amyloid- β (A β) peptide in an AD mouse model. Consistent with these observations, endogenous murine A β generation is also reduced in *Arrb2*^{-/-} mice. Elucidation of the mechanism of the β -arrestin 2-mediated effect on A β levels indicates that recruitment of β -arrestin 2 to two GPCRs implicated in the pathogenesis of AD, GPR3 and the β_2 -adrenergic receptor (β_2 -AR), is required for the promotion of A β release. Collectively, these studies identify β -arrestin 2 as a novel avenue for targeting amyloid pathology and GPCR dysfunction in AD.

Authors' details

¹VIB Center for the Biology of Disease, Leuven, Belgium. ²Center for Human Genetics, University of Leuven (KU Leuven), Leuven, Belgium. ³Neuroscience Medicinal Chemistry, Janssen Research and Development, a Division of Janssen Pharmaceutica NV, Beerse, Belgium.

¹VIB Center for the Biology of Disease, Leuven, Belgium
Full list of author information is available at the end of the article

Published: 13 September 2013

doi:10.1186/1750-1326-8-S1-P41

Cite this article as: Thathiah et al.: β -arrestin 2 regulates A β generation and γ -secretase activity in Alzheimer's disease. *Molecular Neurodegeneration* 2013 **8**(Suppl 1):P41.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

